

k -Limited Erasing Performed by Regular-Controlled Context-Free Grammars

Petr Zemek

(supervised by prof. Alexander Meduna)

Faculty of Information Technology
Brno University of Technology

2010-04-29



Outline

- 1 Light introduction to regulated rewriting
- 2 Motivation and problem description
- 3 Results



From Context-Free Grammars...

Consider the following context-free grammar:

$$S \rightarrow ABC$$

$$A \rightarrow aA$$

$$B \rightarrow bB$$

$$C \rightarrow cC$$

$$A \rightarrow \varepsilon$$

$$B \rightarrow \varepsilon$$

$$C \rightarrow \varepsilon$$



From Context-Free Grammars...

Consider the following context-free grammar:

$$1: S \rightarrow ABC$$

$$2: A \rightarrow aA$$

$$3: B \rightarrow bB$$

$$4: C \rightarrow cC$$

$$5: A \rightarrow \varepsilon$$

$$6: B \rightarrow \varepsilon$$

$$7: C \rightarrow \varepsilon$$



From Context-Free Grammars...

Consider the following context-free grammar:

$$1: S \rightarrow ABC$$

$$2: A \rightarrow aA$$

$$3: B \rightarrow bB$$

$$4: C \rightarrow cC$$

$$5: A \rightarrow \varepsilon$$

$$6: B \rightarrow \varepsilon$$

$$7: C \rightarrow \varepsilon$$



... To Regulated Grammars

Consider the following **regular-controlled** context-free grammar:

$$1: S \rightarrow ABC$$

$$2: A \rightarrow aA$$

$$3: B \rightarrow bB$$

$$4: C \rightarrow cC$$

$$5: A \rightarrow \varepsilon$$

$$6: B \rightarrow \varepsilon$$

$$7: C \rightarrow \varepsilon$$

$$R = \{1\}\{234\}^*\{567\}$$



... To Regulated Grammars

Consider the following regular-controlled context-free grammar:

$$1: S \rightarrow ABC$$

$$2: A \rightarrow aA$$

$$3: B \rightarrow bB$$

$$4: C \rightarrow cC$$

$$5: A \rightarrow \varepsilon$$

$$6: B \rightarrow \varepsilon$$

$$7: C \rightarrow \varepsilon$$

$$R = \{1\}\{234\}^*\{567\}$$

S



... To Regulated Grammars

Consider the following regular-controlled context-free grammar:

$$1: S \rightarrow ABC$$

$$2: A \rightarrow aA$$

$$3: B \rightarrow bB$$

$$4: C \rightarrow cC$$

$$5: A \rightarrow \varepsilon$$

$$6: B \rightarrow \varepsilon$$

$$7: C \rightarrow \varepsilon$$

$$R = \{1\}\{234\}^*\{567\}$$

S



... To Regulated Grammars

Consider the following regular-controlled context-free grammar:

$$1: S \rightarrow ABC$$

$$2: A \rightarrow aA$$

$$3: B \rightarrow bB$$

$$4: C \rightarrow cC$$

$$5: A \rightarrow \varepsilon$$

$$6: B \rightarrow \varepsilon$$

$$7: C \rightarrow \varepsilon$$

$$R = \{1\}\{234\}^*\{567\}$$

$$S \Rightarrow ABC$$



... To Regulated Grammars

Consider the following regular-controlled context-free grammar:

$$1: S \rightarrow ABC$$

$$2: A \rightarrow aA$$

$$3: B \rightarrow bB$$

$$4: C \rightarrow cC$$

$$5: A \rightarrow \varepsilon$$

$$6: B \rightarrow \varepsilon$$

$$7: C \rightarrow \varepsilon$$

$$R = \{1\}\{234\}^*\{567\}$$

$$S \Rightarrow ABC$$



... To Regulated Grammars

Consider the following regular-controlled context-free grammar:

$$1: S \rightarrow ABC$$

$$2: A \rightarrow aA$$

$$3: B \rightarrow bB$$

$$4: C \rightarrow cC$$

$$5: A \rightarrow \varepsilon$$

$$6: B \rightarrow \varepsilon$$

$$7: C \rightarrow \varepsilon$$

$$R = \{1\}\{234\}^*\{567\}$$

$$S \Rightarrow ABC \Rightarrow aABC$$



... To Regulated Grammars

Consider the following regular-controlled context-free grammar:

$$1: S \rightarrow ABC$$

$$2: A \rightarrow aA$$

$$3: B \rightarrow bB$$

$$4: C \rightarrow cC$$

$$5: A \rightarrow \varepsilon$$

$$6: B \rightarrow \varepsilon$$

$$7: C \rightarrow \varepsilon$$

$$R = \{1\}\{234\}^*\{567\}$$

$$S \Rightarrow ABC \Rightarrow aABC$$



... To Regulated Grammars

Consider the following regular-controlled context-free grammar:

$$1: S \rightarrow ABC$$

$$2: A \rightarrow aA$$

$$3: B \rightarrow bB$$

$$4: C \rightarrow cC$$

$$5: A \rightarrow \varepsilon$$

$$6: B \rightarrow \varepsilon$$

$$7: C \rightarrow \varepsilon$$

$$R = \{1\}\{234\}^*\{567\}$$

$$S \Rightarrow ABC \Rightarrow aABC \Rightarrow aAbBC$$



... To Regulated Grammars

Consider the following regular-controlled context-free grammar:

$$1: S \rightarrow ABC$$

$$2: A \rightarrow aA$$

$$3: B \rightarrow bB$$

$$4: C \rightarrow cC$$

$$5: A \rightarrow \varepsilon$$

$$6: B \rightarrow \varepsilon$$

$$7: C \rightarrow \varepsilon$$

$$R = \{1\}\{234\}^*\{567\}$$

$$S \Rightarrow ABC \Rightarrow aABC \Rightarrow aAbBC$$



... To Regulated Grammars

Consider the following regular-controlled context-free grammar:

$$1: S \rightarrow ABC$$

$$2: A \rightarrow aA$$

$$3: B \rightarrow bB$$

$$4: C \rightarrow cC$$

$$5: A \rightarrow \varepsilon$$

$$6: B \rightarrow \varepsilon$$

$$7: C \rightarrow \varepsilon$$

$$R = \{1\}\{234\}^*\{567\}$$

$$S \Rightarrow ABC \Rightarrow aABC \Rightarrow aAbBC \Rightarrow aAbBcC$$



... To Regulated Grammars

Consider the following regular-controlled context-free grammar:

$$1: S \rightarrow ABC$$

$$2: A \rightarrow aA$$

$$3: B \rightarrow bB$$

$$4: C \rightarrow cC$$

$$5: A \rightarrow \varepsilon$$

$$6: B \rightarrow \varepsilon$$

$$7: C \rightarrow \varepsilon$$

$$R = \{1\}\{234\}^*\{567\}$$

$$S \Rightarrow ABC \Rightarrow aABC \Rightarrow aAbBC \Rightarrow aAbBcC$$



... To Regulated Grammars

Consider the following regular-controlled context-free grammar:

$$1: S \rightarrow ABC$$

$$2: A \rightarrow aA$$

$$3: B \rightarrow bB$$

$$4: C \rightarrow cC$$

$$5: A \rightarrow \varepsilon$$

$$6: B \rightarrow \varepsilon$$

$$7: C \rightarrow \varepsilon$$

$$R = \{1\}\{234\}^*\{567\}$$

$$S \Rightarrow ABC \Rightarrow aABC \Rightarrow aAbBC \Rightarrow aAbBcC \Rightarrow abBcC$$



... To Regulated Grammars

Consider the following regular-controlled context-free grammar:

$$1: S \rightarrow ABC$$

$$2: A \rightarrow aA$$

$$3: B \rightarrow bB$$

$$4: C \rightarrow cC$$

$$5: A \rightarrow \varepsilon$$

$$6: B \rightarrow \varepsilon$$

$$7: C \rightarrow \varepsilon$$

$$R = \{1\}\{234\}^*\{567\}$$

$$S \Rightarrow ABC \Rightarrow aABC \Rightarrow aAbBC \Rightarrow aAbBcC \Rightarrow abBcC$$



... To Regulated Grammars

Consider the following regular-controlled context-free grammar:

$$1: S \rightarrow ABC$$

$$2: A \rightarrow aA$$

$$3: B \rightarrow bB$$

$$4: C \rightarrow cC$$

$$5: A \rightarrow \varepsilon$$

$$6: B \rightarrow \varepsilon$$

$$7: C \rightarrow \varepsilon$$

$$R = \{1\}\{234\}^*\{567\}$$

$$S \Rightarrow ABC \Rightarrow aABC \Rightarrow aAbBC \Rightarrow aAbBcC \Rightarrow abBcC \Rightarrow abcC$$



... To Regulated Grammars

Consider the following regular-controlled context-free grammar:

$$1: S \rightarrow ABC$$

$$2: A \rightarrow aA$$

$$3: B \rightarrow bB$$

$$4: C \rightarrow cC$$

$$5: A \rightarrow \varepsilon$$

$$6: B \rightarrow \varepsilon$$

$$7: C \rightarrow \varepsilon$$

$$R = \{1\}\{234\}^*\{567\}$$

$$S \Rightarrow ABC \Rightarrow aABC \Rightarrow aAbBC \Rightarrow aAbBcC \Rightarrow abBcC \Rightarrow abcC$$



... To Regulated Grammars

Consider the following regular-controlled context-free grammar:

$$1: S \rightarrow ABC$$

$$2: A \rightarrow aA$$

$$3: B \rightarrow bB$$

$$4: C \rightarrow cC$$

$$5: A \rightarrow \varepsilon$$

$$6: B \rightarrow \varepsilon$$

$$7: C \rightarrow \varepsilon$$

$$R = \{1\}\{234\}^*\{567\}$$

$$S \Rightarrow ABC \Rightarrow aABC \Rightarrow aAbBC \Rightarrow aAbBcC \Rightarrow abBcC \Rightarrow abcC \Rightarrow abc$$



... To Regulated Grammars

Consider the following regular-controlled context-free grammar:

$$1: S \rightarrow ABC$$

$$2: A \rightarrow aA$$

$$3: B \rightarrow bB$$

$$4: C \rightarrow cC$$

$$5: A \rightarrow \varepsilon$$

$$6: B \rightarrow \varepsilon$$

$$7: C \rightarrow \varepsilon$$

$$R = \{1\}\{234\}^*\{567\}$$

$$S \Rightarrow ABC \Rightarrow aABC \Rightarrow aAbBC \Rightarrow aAbBcC \Rightarrow abBcC \Rightarrow abcC \Rightarrow abc$$

$$L = \{a^n b^n c^n \mid n \geq 0\}$$



Open Problem

Can any
regular-controlled context-free grammar **with** erasing rules
be transformed into an equivalent
regular-controlled context-free grammar **without** erasing rules?



Main Result

Condition Erasure of nonterminals in a k -limited way, $k \geq 0$.



Main Result

Condition Erasure of nonterminals in a k -limited way, $k \geq 0$.

+

Algorithm Elimination of all erasing rules from any regular-controlled context-free grammar satisfying this condition.



Main Result

Condition Erasure of nonterminals in a k -limited way, $k \geq 0$.

+

Algorithm Elimination of all erasing rules from any regular-controlled context-free grammar satisfying this condition.

=

Partial solution of the problem.



Thank you for your attention!